

**Publikationen:**

1. Arthroscopic repair of HAGL lesions yields good clinical results, but may not allow return to former level of sport.  
**Schmiddem U**, Watson A, Perriman D, Liodakis E, Page R  
Knee Surg Sports Traumatol Arthrosc. 2019 Oct;27(10):3246-3253. doi: 10.1007/s00167-019-05414-5. Epub 2019 Feb 27.
2. Monocortical fixation of the coracoid in the Latarjet procedure is significantly weaker than bicortical fixation.  
**Schmiddem U**, Hawi N, Liodakis E, Dratzidis A, Kraemer M, Hurschler C, Page R, Meller R  
Knee Surg Sports Traumatol Arthrosc. 2018 Jan 13. doi: 10.1007/s00167-018-4837-2
3. Fatty degeneration of the deltoid muscle associated with displaced proximal humerus fractures.  
Panzica M, Brandes J, **Schmiddem U**, Karch A, Hawi N, Krettek C, Meller R  
Technol Health Care. 2017 Oct 23;25(5):959-967. doi: 10.3233/THC-170834.
4. Combined Fluoroscopic and Arthroscopic Detection and Removal of a Foreign Body Lost During Elective Shoulder Arthroscopy: A Case Report.  
**Schmiddem U**, Hawi N, Suero EM, Meller R  
J Orthop Case Rep. 2017 Mar-Apr;7(2):78-81. doi: 10.13107/jocr.2250-0685.760.
5. Arthroscopic Debridement for Irreparable Rotator Cuff Tears.  
N. Hawi, **U. Schmiddem**, M. Omar, T. Stuebig, C. Krettek, M. Petri, R. Meller  
The Open Orthopaedics Journal, 2016, 10: 324-329
6. Posttraumatic nonunions and malunions of the proximal humerus: Possibilities and limitations of corrective osteotomy.  
Meller R, Hawi N, **Schmiddem U**, Millett PJ, Petri M, Krettek C  
Unfallchirurg. 2015 Jun 25. German.
7. Indikationen und Ergebnisse debridierender Eingriffe bei irreparabler Rotatorenmanschettenruptur.  
Hawi N, **Schmiddem U**, Liodakis E, Petri M, Krettek C, Meller R  
Arthroskopie 2014 February; 27: 16 -20
8. Klinische und apparative Untersuchungstechniken des Schultergelenkes.  
OP-Journal 2013; 29(1): 6-14. DOI: 10.1055/s-0032-1328427  
**Schmiddem U**, Krettek C, Meller R
9. First degree open, proximal and distal fractures of the lower arm with scapholunate ligament rupture. Case report of a severe extremity trauma.  
**Schmiddem U**, Jenzer A.  
Unfallchirurg. 2013 Dec;116(12):1128-32. doi: 10.1007/s00113-012-2313-x. German.
10. FTY720 for treatment of ischemia reperfusion injury following complete renal ischemia – impact on long term survival and T-lymphocyt tissue infiltration.  
Kaudel CP, Frink M, **Schmiddem U**, Probst C, Bergmann S, Krettek C, Klempnauer J, van Griensven M, Winkler M  
Transplant Proc. 2007 Mar;39(2): 499-502
11. FTY720 application following isolated warm liver ischemia improves long term survival and organ protection in a mouse model.  
Kaudel CP, Frink M, van Griensven M, **Schmiddem U**, Probst C, Bergmann S, Krettek C, Klempnauer J, Winkler M  
Transplant Proc. 2007 Mar;39(2): 493-8

12. FTY720 for treatment of ischemia reperfusion injury following complete renal ischemia in C57/BL6 mice.  
Kaudel CP, **Schmiddem U**, Frink M, Bergmann S, Pape HC, Krettek C, Klempnauer J, Winkler M  
Transplant Proc. 2006 Apr;38(3): p. 679-81.

*U. Schmid*